Founding Mothers of Jewish Communities: Geographically Separated Jewish Groups Were Independently Founded by Very Few Female Ancestors

Mark G. Thomas,^{1,*} Michael E. Weale,^{1,*} Abigail L. Jones,¹ Martin Richards,³ Alice Smith,² Nicola Redhead,² Antonio Torroni,^{5,6}






Table 1

Y Chromosome Diversity within 18 Jewish and Non-Jewish Populations


en la conserva de la

Table 2

mtDNA Diversity within 18 Jewish and Non-Jewish Populations

	•••	1.1.1.1	• • ₆ • _K	
-• n h ()	- h (п	h (• • • • •

Table 3 Frequently Encountered mtDNA Haplotypes



، به محمد محمد محمد به محمد محمد محمد محمد به م محمد به ď ?


Table 4

Mean $F_{\rm ST}$ Values in mtDNA and Y Chromosome Data

	■/ I+ f		
()	1 ? /	<i>, , , , , , , , , ,</i>	
I I (⊉1) I ● (1) ● ● (1) ●			í í í

(P < 0) = (P <

 $\mathbf{c} = \mathbf{c} + \mathbf{c} +$



Discussion







Table 5

Comparison of X Chromosome LD among Georgian Jews and non-Jewish British Men

••						
k /	*	\$	≤	≤	\$	
$ \begin{array}{c} P \leqslant \\ \end{array} $	• 1 1	2	2	1	1	

 $\mathbf{r}_{\mathbf{r}_{i}} = \mathbf{r}_{i} \mathbf{r}_{\mathbf{r}_{i}} + \mathbf{r}_{i} \mathbf$



Am. J. Hum. Genet. 70:1411-1420, 2002

#